

WHAT RISK GROUP ARE YOU?

ARE YOU AT AVERAGE-RISK?

People 45 or older:

- Without prior colorectal cancer or polyps
- Without any of the factors that define high-risk screening

EVERY

YEAR

TEST: EVERY 10 YRS Colonoscopy

TEST:

Fecal Immunohistochemistry (Stool FIT) TEST:

Multi-Target Stool DNA (mt-sDNA) EVERY 3^{YRS}

Colonoscopy negative result:

The test only needs to be repeated every 10 years.

FIT or mt-sDNA screening postive result:

A colonoscopy to find the suspected cancer and locate and remove cancerous polyps will be required.

DO YOU HAVE SYMPTOMS?

- · Rectal bleeding
- · Anemia
- · Change in bowel habits
- Persistent abdominal pain
- Unintentional weight loss

TEST: Colonoscopy DO YOU HAVE A PERSONAL HISTORY?

- Previously removed pre-cancerous colorectal polyps
- Previously had colorectal cancer

TEST: Colonoscopy

ARI HIG

ARE YOU AT HIGH-RISK?

- Family history of colorectal cancer or precancerous polyps in a first degree relative diagnosed before age 60
- Multiple first-degree relatives with colorectal cancer or precancerous polyps
- · Family history of inherited colorectal cancer syndrome
- Previous diagnosis of ulcerative colitis or Crohn's disease

TEST: Colonoscopy

Screening recommended
BEFORE 45 YEARS

if you have symptoms, personal history or are at high risk.

SURVIVAL RATE
when detected early.

1in3
U.S. ADULTS
are not getting screened for

colon cancer as recommended.2

Nearly

Learn more at ASGE.org/Screening



©2022 ASGE. All Rights Reserved.

