

The role of endoscopy in the management of constipation

This is one of a series of statements discussing the use of GI endoscopy in common clinical situations. The Standards of Practice Committee of the American Society for Gastrointestinal Endoscopy (ASGE) prepared this guideline. In preparing this document, MEDLINE databases were used to search for publications pertaining to this topic between January 1990 and December 2013. Additional references were obtained from the bibliographies of the identified articles and from recommendations of expert consultants. When few or no data exist from well-designed, prospective trials, emphasis was given to results from large series and reports from recognized experts. The reported evidence and recommendations on the basis of reviewed studies were based on consensus opinion of the strength of the supporting evidence (Table 1).¹ The strength of individual recommendations is based on both the aggregate evidence quality and an assessment of the anticipated benefits and harms. Weaker recommendations are indicated by phrases such as “We suggest...,” whereas stronger recommendations are typically stated as “We recommend...”

ASGE guidelines for the appropriate use of endoscopy are based on a critical review of the available data and expert consensus at the time that the documents are drafted. Further controlled clinical studies may be needed to clarify aspects of this document. This document may be revised as necessary to account for changes in technology, new data, or other aspects of clinical practice and is solely intended to be an educational device to provide information that may assist endoscopists in providing care to patients. This document is not a rule and should not be construed as establishing a legal standard of care or as encouraging, advocating, requiring, or discouraging any particular treatment. Clinical decisions in any particular case involve a complex analysis of the patient's condition and available courses of action. Therefore, clinical considerations may lead an endoscopist to take a course of action that varies from the recommendations and suggestions proposed in this document.

INTRODUCTION AND EPIDEMIOLOGY

Constipation is a common symptom affecting 2% to 27% of the population and resulting in about 2.5 million

physician visits in the United States annually.^{2,3} The prevalence of constipation is higher in women than in men⁴ and increases with age.⁵ Low socioeconomic status, physical inactivity, a history of sexual abuse, and depression have all been reported to be risk factors for constipation.⁶

DEFINITION

Chronic constipation has been defined by the Rome III diagnostic criteria (Table 2).⁷ Constipation symptoms include excessive straining, discomfort at defecation, or passage of hard or pellet-like stools, even though the frequency of defecation may be normal.

THE ROLE OF ENDOSCOPY

Patients with constipation should undergo colonoscopy if they have rectal bleeding, heme-positive stool, iron deficiency anemia, weight loss, or obstructive symptoms. In addition, colonoscopy should be considered in selected patients to exclude obstruction from cancer, stricture, and extrinsic compression. Colonoscopy also should be done prior to surgery for constipation. In younger patients, a flexible sigmoidoscopy may be sufficient to exclude distal disease. Suspected Hirschsprung's disease requires anorectal manometry and deep biopsy to examine for the absence of myenteric neurons.⁸⁻¹⁰

Patients aged >50 years who have not had prior colorectal cancer screening should undergo colonoscopy. Studies evaluating the association of chronic constipation and colorectal cancer have produced inconsistent findings. Chronic constipation was associated with an increased risk of colon cancer in two U.S. population-based retrospective studies^{11,12} but not in a prospective study of female nurses.¹³ A retrospective study from Australia also reported increased cancer risk in patients with constipation,¹⁴ and a retrospective study from Japan found increased risk in those who used laxatives frequently.¹⁵ However, a meta-analysis of 28 studies (8 cross-sectional surveys, 3 cohort studies, 17 case-control studies) demonstrated no increase in colorectal cancer in patients with chronic constipation.¹⁶

The yield of colonoscopy in isolated constipation is low and comparable to that of asymptomatic patients undergoing colonoscopy for colorectal cancer screening. In one study of 563 sigmoidoscopies or colonoscopies done for the evaluation of constipation, colorectal cancer was found in 8 (1.4%), adenomas in 82 (14.6%), and advanced

TABLE 1. GRADE system[†] for rating the quality of evidence for guidelines[†]

Quality of evidence	Definition	Symbol
High quality	Further research is very unlikely to change our confidence in the estimate of effect.	⊕⊕⊕⊕
Moderate quality	Further research is likely to have an important impact on our confidence in the estimate of effect and may change the estimate.	⊕⊕⊕○
Low quality	Further research is very likely to have an important impact on our confidence in the estimate of effect and is likely to change the estimate.	⊕⊕○○
Very low quality	Any estimate of effect is very uncertain.	⊕○○○

GRADE, Grading of Recommendations Assessment, Development, and Evaluation.

[†]Adapted with permission from Guyatt et al.¹

TABLE 2. Rome III criteria for functional constipation[†]

Criteria fulfilled for the previous 3 months, with symptom onset ≥ 6 months prior to diagnosis:

1. Must include ≥ 2 of the following:

- a. Straining during at least 25% of defecations
- b. Lumpy or hard stools in at least 25% of defecations
- c. Sensation of incomplete evacuation for at least 25% of defecations
- d. Sensation of anorectal obstruction/blockage for at least 25% of defecations
- e. Manual maneuvers to facilitate at least 25% of defecations

f. Fewer than 3 defecations per week

2. Loose stools are rarely present without the use of laxatives.

3. Insufficient criteria for irritable bowel syndrome.

lesions (cancer or adenoma with malignancy, high-grade dysplasia, villous features, or size >10 mm) in 24 (4.3%).¹⁷ Another study that evaluated the yield of colonoscopy performed for the sole indication of constipation found that the prevalence of colorectal neoplasia was lower in patients with constipation than in those undergoing colonoscopy for routine colorectal cancer screening.¹⁸ A retrospective review of 41,775 index colonoscopies performed for colorectal cancer screening, constipation alone, or constipation with another indication found that patients with constipation alone had a lower risk of significant findings than patients undergoing colonoscopy for average-risk screening.¹⁹ Associated findings may include solitary rectal ulcer syndrome (indicating rectal prolapse), anal fissure, and melanosis coli (indicating chronic laxative use).

Colonoscopy may be used to provide therapy in some patients. Fibrotic strictures from inflammatory bowel disease,

surgery, or ischemia can be dilated at the time of colonoscopy.²⁰⁻²³ Percutaneous endoscopic cecostomy or colostomy has been used with favorable results in children with severe refractory constipation caused by conditions such as neurogenic bowel.^{24,25} In adults with acute colonic pseudo-obstruction and neurogenic bowel, percutaneous endoscopic cecostomy may be effective when conservative treatment fails.²⁶ It is important to understand that colonoscopy has no role in stool disimpaction, although there are reports of colonoscopic removal of bezoar-induced fecal impaction.²⁷

Chronic constipation is an independent risk factor for inadequate bowel preparation for colonoscopy.²⁸ In these patients, a more aggressive regimen for colon cleansing should be considered.

SUMMARY

1. We recommend that GI endoscopy should not be performed in the initial evaluation of patients presenting with symptoms of chronic constipation in the absence of alarm features or suspicion of organic GI disease. ⊕⊕⊕○
2. We recommend that patients with constipation undergo colonoscopy to exclude organic disease if they have rectal bleeding, heme-positive stool, iron deficiency anemia, or weight loss prior to surgical therapy for chronic constipation. ⊕⊕⊕⊕
3. We recommend that patients aged >50 years presenting with constipation who have not previously had colon cancer screening should have a colonoscopy. ⊕⊕⊕⊕
4. We recommend colonoscopy to allow dilation of benign colon strictures and creation of percutaneous cecostomy when clinically appropriate and feasible. ⊕⊕⊕○

DISCLOSURES

M. Khashab is a consultant for Boston Scientific and Olympus America and has received research support

from Cook Medical. B. Cash is a consultant and member of the speakers bureaus for Takeda, Forest and Ironwood Pharmaceuticals. All other authors disclosed no financial relationships relevant to this article.

REFERENCES

- Guyatt G, Oxman AD, Akl EA, et al. GRADE guidelines: introduction- GRADE evidence profiles and summary of findings tables. *J Clin Epidemiol* 2011;64:383-94.
- Sonnenberg A, Koch TR. Physician visits in the United States for constipation: 1958-1986. *Dig Dis Sci* 1989;34:606-11.
- Pare P, Ferrazzi S, Thompson WG, et al. An epidemiological survey of constipation in Canada: definitions, rates, demographics, and predictors of health care seeking. *Am J Gastroenterol* 2001;96:3130-7.
- Heaton KW, Radvan J, Cripps H, et al. Defecation frequency and timing, and stool form in the general population: a prospective study. *Gut* 1992;33:818-24.
- Johanson JF, Sonnenberg A, Koch TR. Clinical epidemiology of chronic constipation. *J Clin Gastroenterol* 1989;11:525-36.
- Everhart JE, Go VL, Johannes RS, et al. A longitudinal survey of self-reported bowel habits in the United States. *Dig Dis Sci* 1989;34:1153-62.
- Longstreth G, Thompson WG, Chey WD, et al. Functional bowel disorders. *Gastroenterology* 2006;130:1480-91.
- Wald A. Approach to the patient with constipation. In: Yamada T, editor. *Textbook of Gastroenterology*, 4th ed. Philadelphia: Lippincott, Williams & Wilkins; 2003. p. 894-910.
- Taxman TL, Yulish BS, Rothstein FC. How useful is the barium enema in the diagnosis of infantile Hirschsprung's disease? *Am J Dis Child* 1986;140:881-4.
- Aldridge RT, Campbell PE. Ganglion cells distribution in the normal rectum and anal canal. A basis for diagnosis of Hirschsprung's disease by anorectal biopsy. *J Pediatr Surg* 1968;3:475-89.
- Roberts MC, Millikan RC, Galanko JA, et al. Constipation, laxative use, and colon cancer in a North Carolina population. *Am J Gastroenterol* 2003;98:857-64.
- Jacobs EJ, White E. Constipation, laxative use, and colon cancer among middle-aged adults. *Epidemiology* 1998;9:385-91.
- Dukas L, Willett WC, Colditz GA, et al. Prospective study of bowel movement, laxative use, and the risk of colorectal cancer among women. *Am J Epidemiol* 2000;151:958-64.
- Kune GA, Kune S, Field B, et al. The role of chronic constipation, diarrhea, and laxative use in the etiology of large-bowel cancer. Data from the Melbourne Colorectal Cancer Study. *Dis Colon Rectum* 1988;31:507-12.
- Watanabe T, Nakaya N, Kurashima K, et al. Constipation, laxative use and risk of colorectal cancer: the Miyagi Cohort Study. *Eur J Cancer* 2004;40:2109-15.
- Power AM, Talley NJ, Ford AC. Association between constipation and colorectal cancer: systematic review and meta-analysis. *Am J Gastroenterol* 2013;108:894-903.
- Pepin C, Ladabaum U. The yield of lower endoscopy in patients with constipation: survey of a university hospital, a public county hospital and a veterans administration medical center. *Gastrointest Endosc* 2002;56:325-32.
- Obusez EC, Lian L, Kariv R, et al. Diagnostic yield of colonoscopy for constipation as the sole indication. *Colorectal Dis* 2012;14:589-91.
- Gupta M, Holub J, Knigge K, et al. Constipation is not associated with an increased risk of findings on colonoscopy: results from a national endoscopy consortium. *Endoscopy* 2010;43:208-12.
- Virgilio C, Cosentino S, Favara C, et al. Endoscopic treatment of post-operative colonic strictures using an achalasia dilator: short-term and long-term results. *Endoscopy* 1995;27:219-22.
- Truong S, Willis S, Schumpelick V. Endoscopic therapy of benign anastomotic strictures of the colorectum by electroincision and balloon dilatation. *Endoscopy* 1997;29:845-9.
- Sabate JM, Villarejo J, Bouhnik Y, et al. Hydrostatic balloon dilatation of Crohn's strictures. *Aliment Pharmacol Ther* 2003;18:409-13.
- Morini S, Hassan C, Lorenzetti R, et al. Long-term outcome of endoscopic pneumatic dilatation in Crohn's disease. *Dig Liver Dis* 2003;35:893-7.
- Rawat DJ, Haddad M, Geoghegan N, et al. Percutaneous endoscopic colostomy of the left colon: a new technique for management of intractable constipation in children. *Gastrointest Endosc* 2004;60:39-43.
- Rivera MT, Kugathasan S, Berger W, et al. Percutaneous colonoscopic cecostomy for management of chronic constipation in children. *Gastrointest Endosc* 2001;53:225-8.
- Ramage JI Jr, Baron TH. Percutaneous endoscopic cecostomy: a case series. *Gastrointest Endosc* 2003;57:752-5.
- Purcell L, Gremse DA. Sunflower seed bezoar leading to fecal impaction. *South Med J* 1995;88:87-8.
- Ness RM, Manam R, Hoen H, et al. Predictors of inadequate bowel preparation for colonoscopy. *Am J Gastroenterol* 2001;96:1797-802.

Prepared by:

ASGE STANDARDS OF PRACTICE COMMITTEE

Brooks D. Cash, MD, Committee Chair

Ruben D. Acosta, MD

Vinay Chandrasekhara, MD

Krishnavel V. Chathadi, MD

Mohammad A. Eloubeidi, MD

Robert D. Fanelli, MD, SAGES Representative

Ashley L. Faulx, MD

Lisa Fonkalsrud, RN, BSN, CGRN, SGNA Representative

Mouen A. Khashab, MD

Jenifer R. Lightdale, MD, MPH

V. Raman Muthusamy, MD

Shabana F. Pasha, MD

John R. Saltzman, MD

Aasma Shaukat, MD, MPH

Amy Wang, MD

This document is a product of the Standards of Practice Committee. This document was reviewed and approved by the Governing Board of the American Society for Gastrointestinal Endoscopy.